

Eloisa Fornaro Clemente^{1,2,3}, Mariana Prezzotto Pereira^{2,3,4}, Livia Maria Pereira da Silva^{2,3,4}, Giovana de Freitas^{1,2,3}, Joao Pessoa Araújo Jr^{2,5}, Bruno Cesar Rossini^{2,5}, Bruna Cavecci Mendonça^{1,2,3,6}, Lucilene Delazari dos Santos^{1,2}

¹. FMB/UNESP, Faculdade de Medicina de Botucatu, Universidade Estadual Paulista "Júlio de Mesquita Filho", Av. Prof. Montenegro, s/n - Distrito de Rubião Jr, Botucatu - SP, 18618-687;

². IBTEC/UNESP, Instituto de Biotecnologia, Universidade Estadual Paulista "Júlio de Mesquita Filho", Alameda das Tecomarias, s/n - Chácara Capão Bonito - Botucatu/SP - 18607-440;

³. Triad for Life Ltda, Paulo Daniel Leite Pesquisa e Desenvolvimento, Incubadora Tecnológica Prospecta, Rua Doutor Jose Barbosa de Barros, 1780, SALA 6, Jardim Paraíso, Botucatu/SP - 18610-307;

⁴. FCA/UNESP, Faculdade de Ciências Agrônômicas, Universidade Estadual Paulista "Júlio de Mesquita Filho", Av. Universitária, nº 3780 - Altos do Paraíso, Botucatu - SP, 18610-034;

⁵. IBB/UNESP, Instituto de Biociências, Universidade Estadual Paulista "Júlio de Mesquita Filho", Rua Prof. Dr. Antônio Celso Wagner Zanin, 250 - Distrito de Rubião Junior - Botucatu/SP - 18618-689;

⁶. CEVAP/UNESP, Centro de Estudos de Venenos e Animais Peçonhentos da Universidade Estadual Paulista "Júlio de Mesquita Filho", Rodovia Alcides Soares, Km 3, Portaria 2 - Fazenda Experimental Lageado, UNESP - Botucatu -SP/SP - CEP 18619-002;

Snakebite accidents are a potentially fatal medical emergency resulting from snakebite envenomation. Considered by the World Health Organization (WHO) as a neglected tropical disease, snakebites affect approximately 28,000 Brazilians annually. In Brazil, *Bothrops* genus snakes are the most epidemiologically significant, and the pathophysiology of these incidents includes proteolytic, coagulant, and hemorrhagic mechanisms, with local and/or systemic manifestations. The unique effective treatment is serotherapy, which can lead to adverse effects such as anaphylactic reactions and serum sickness, related to the manufacturing practices and purity of the antivenom. The aim of this study was to evaluate the efficacy of coupling the *Bothrops jararaca* snake venom to an affinity chromatography matrix and to identify the proteins of a commercial *Bothrops* antivenom that specifically interact with the toxins of this snake's venom. As this study is part of a public-private research and development agreement, the methodology cannot be detailed. In summary, low-pressure liquid chromatography using the AKTA Pure equipment (Cytiva) and shotgun mass spectrometry using the Q-Exactive mass spectrometry equipment (Thermo Fisher) were employed in this study. As a result, the affinity chromatography matrix prepared was able to purify the antivenom fraction that has greater specificity with the toxins of the *Bothrops* venom. The purified proteins were characterized and 68% of them were classified as therapeutically relevant for the treatment of a snakebite (immunoglobulins), and 10% of albumin. On the other hand, analyses of the crude antivenom showed 45% immunoglobulins and 27% albumin. Additionally, the characterization of the snake venom profile coupled to the chromatography matrix revealed 100% of the immunogenic protein classes of *Bothrops* venom, compared to the previously characterized crude venom. However, the only class of non-coupled toxins were bradykinin-potentiating peptides, which do not have immunological potential. This proposal demonstrates that this affinity chromatography approach is effective in obtaining specific antibodies present in bothropic antivenoms, and these findings can support further studies by production centers regarding the enrichment of commercial snake antivenoms.

Agradecimentos: The authors appreciate the valuable financial support by PIBIC/UNESP; CNPq (315026/2021-9); PIPE FAPESP (2018/15446-8); PIPE FAPESP (2020/09819-6); PIPE FAPESP (2022/02869-3); FAPESP (2019/14647-2; 2020/16747-1; 2021/03671-0; 2022/13458-4; 2023/00804-4).